

Measuring Star-Formation Histories at $z \sim 2$ with a Semi-Resolved Approach


Sam Cutler, Kate Whitaker, Mauro Giavalisco, Zhiyuan Ji, Yingjie Cheng
UMass Amherst, DAWN-IRES Senior Scholar/Cosmic Dawn Center
@secutler on Twitter

2014-02-05


Orbits: 59

Pointings: 44

Credit: Iva Momcheva,
3D-DASH Team






Credit: Sune Toft






Semi-resolved Prospector SEDs of star-forming galaxies

Does SFH differ between galactic centers and outskirts?





Semi-resolved Prospector SEDs of star-forming galaxies

Does SFH differ between galactic centers and outskirts?





Semi-resolved Prospector SEDs of star-forming galaxies

Centers of SFGs form fast and late compared to the outskirts




Semi-resolved Prospector SEDs of star-forming galaxies

Galaxies still on the main sequence after centers form




Timescales of quenched galaxies with color gradients

Compare (central) quenching and (global) formation timescales




Data from REQUIEM (Akhshik+23)





Timescales of quenched galaxies with color gradients

Study timescales with a larger sample (e.g. 3D-DASH)






+





Timescales of quenched galaxies with color gradients

Use REQUIEM SFHs to create grid of model color-gradients



Timescales of quenched galaxies with color gradients

Use REQUIEM SFHs to create grid of model color-gradients






DAWN



Cutler+ in prep.

Timescales of quenched galaxies with color gradients




Use observed i - H colors to approximate timescales of $z \sim 2$ galaxies in 3D-DASH imaging



Cutler+ in prep.

Timescales of quenched galaxies with color gradients

Use observed *i-H* colors to approximate timescales of $z \sim 2$ galaxies in 3D-DASH imaging



Cutler+ in prep.

Summarizing...


(arXiv:2208.01653)




1. **Cutler et al. 2023:** *Prospector SFHs of centers/outskirts of star-forming galaxies*
 - For $\log M < 10.5$, centers form later in a rapid burst of star-formation
 - Galaxies remain on the main sequence even after central starburst (not quenching yet)
2. **Cutler et al. in prep:** *Estimate formation & quenching timescales of quiescent galaxies*
 - Measure observed $i-H$ color gradients from COSMOS/3D-DASH and predict timescales
 - Quenched 3D-DASH galaxies follow a similar trend in $t_{50} - \tau_{\text{center}}$ as REQUIEM



Contact me with any thoughts or suggestions!


secutler@umass.edu; [@secutler](#) on Twitter; samecutler.github.io






Bonus Slides!

























Center Covariance (ID=12024)



Outskirts Covariance (ID=12024)



Total Covariance (ID=12024)

